



ANTI-INFLAMMATORY EFFECT OF METHANOLIC EXTRACT OF *TAMILNADIA ULIGINOSA* (RETZ.) ROOTS

Dorababu Neerugatti*, Ganga Rao Battu, Ravi Kumar Jangiti

A.U. College of Pharmaceutical Sciences, Andhra University, Visakhapatnam, A.P, India-530003.

ABSTRACT

To evaluate the anti-inflammatory activity of the roots of *Tamilnadia uliginosa*. Anti-inflammatory activity was conducted by using Carrageenan induced rat paw oedema model. The thickness of paws of rats was measured every hour up to 5 hrs for methanolic extract at doses of 100, 200 and 400mg/kg by volume displacement method, using plethysmometer. All doses of the extract (methanol) from the roots of *Tamilnadia uliginosa* showed significant percentage of inhibition on Carrageenan induced paw oedema; methanol extract from the roots of *Tamilnadia uliginosa* at dose 400mg/kg produced significant ($P < 0.001$) percentage of inhibition when compared to other doses. *Tamilnadia uliginosa* roots extract (methanol), showed significant anti-inflammatory activity.

Key words: *Tamilnadia uliginosa*, Roots, Anti-inflammatory activity.

INTRODUCTION

Inflammation is the response to injury of cells and body tissues through different factors such as infections, chemicals, and thermal and mechanical injuries (Oyedapo *et al.*, 2008). At the very early stage of inflammation, neutrophils are the first cells that migrate to the inflammatory sites under the regulation of molecules produced by rapidly responding macrophages and mast cells predisposed in tissues (Coussens, 2002; Nathan, 2002). As the inflammation progresses, various types of leukocytes, lymphocytes, and other inflammatory cells are activated and attracted to the inflamed site by a signaling network involving a great number of growth factors, cytokines and chemokines (Coussens, 2002; Nathan, 2002). Nonsteroidal anti-inflammatory drugs (NSAIDs) are the most clinically important medicine used for the treatment of inflammation-related diseases like arthritis, asthma, and cardiovascular disease (Conforti *et al.*, 2009). Nonsteroidal anti-inflammatory drugs (NSAIDs) are among the most widely used medications due to their efficacy for a wide range of pain and inflammatory

conditions (Jacob Jesurun *et al.*, 2013). However, the long-term administration of NSAID may induce gastrointestinal ulcers, bleeding, and renal disorders due to their nonselective inhibition of both constitutive (COX-1) and inducible (COX-2) isoforms of the cyclooxygenase enzymes (Robert, 1976; Peskar, 1977; Tapiero *et al.*, 2002). Therefore, new anti-inflammatory and analgesic drugs lacking those effects are being searched all over the world as alternatives to NSAIDs and opiates (Dharmasiri *et al.*, 2003; Kumara, 2001). Medicinal plants are believed to be an important source of new chemical substances with potential therapeutic effects. The research into plants with alleged folkloric use as pain relievers, anti-inflammatory agents, should therefore be viewed as a fruitful and logical research strategy in the search for new analgesic and anti-inflammatory drugs (Gupta *et al.*, 2006). In the present study different doses of methanolic extract of *Tamilnadia uliginosa* roots were investigated for anti-inflammatory activity.

MATERIALS AND METHODS

Preparation of extracts from the roots of *Tamilnadia uliginosa*

The roots of *Tamilnadia uliginosa* were collected from Tirumala hills, Chittoor, Andhra Pradesh, India and authenticated by Dr. Madhava shetty,

Corresponding Author

Dorababu Neerugatti

Email: ndorababu81@gmail.com

taxonomist, Department of Botany, Sri Venkateswara University, Andhra Pradesh. Shade dried roots of *Tamilnadia uliginosa* were powdered and separately extracted in a Soxhlet apparatus for 6 hrs with methanol and then concentrated under vacuum at temperature of 45°C by using rotary evaporator, dried completely, weighed and stored in desiccators.

Drug and Chemicals

Ibuprofen, Carboxy methyl cellulose sodium (Na CMC) and Carrageenan were purchased from Sigma chemicals, USA. All chemicals used in the study were of analytical grade.

Animals

Albino rats (National Institute of Nutrition, Hyderabad, India) of either sex weighing 200-250gm were used in the studies. The animals were maintained under standard laboratory conditions at an ambient temperature of 23±2°C having 50±5% relative humidity with 12h light and dark cycle. The use and care of the animals in the experimental protocol has been approved by the local Institutional Animal Ethics Committee (Regd. No. 516/01/A/CPCSEA) following the guidelines of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), Government of India.

Acute inflammation model: carrageenan induced rat paw Oedema

Inflammation was induced in the left hind paw of each rat by sub-plantar injection of 1% carrageenan suspension (0.1ml). The right hind paw of the rat was injected 0.1ml of saline. Group-I received drug vehicle (1% sodium CMC). Group-II received standard drug

Ibuprofen at a dose of 10 mg/kg b.w. Group-III, IV and V received orally methanolic extracts of *Tamilnadia uliginosa* roots at doses of 100mg/kg 200mg/kg and 400mg/kg b.w in sodium carboxy methyl cellulose suspension one hour prior to the induction of oedema by carrageenan injection. The paw volume of rats was measured at 0 hr and hourly intervals up to 5th hour after the administration of carrageenan by volume displacement method, using plethysmometer. (Duffy et al., 2002)

The percent inhibition of paw thickness was calculated using the formula:

$$\text{Percentage inhibition} = 100[1 - (Y_t / Y_c)]$$

Y_t = Average increase in thickness of paw in groups tested with test compounds.

Y_c = Average increase in thickness of paw in control

Statistical Analysis

Data of Paw thickness was analyzed by using One-Way ANOVA followed by post hoc test Dunnett's test using Graph pad Prism-5 software. The results were expressed as Mean ±S.E.M. P<0.05 was considered as significant.

RESULTS

Carrageenan induced rat paw Oedema

Tamilnadia uliginosa roots extracts at 100 mg/kg, 200 mg/kg and 400mg/kg doses and Ibuprofen (10 mg/kg b.w.) inhibited the maximal paw oedema response by 17, 24, 57 and 72 respectively during the 5 h of the carrageenan-induced rat paw acute inflammation. Volume of mercury displaced in ml at various time intervals in hours is recorded. The results of were shown in Table 1 and 2. Figure1.


Table 1. Anti-inflammatory activity of *Tamilnadia uliginosa* extracts on carrageenan induced paw oedema in rats

S. No	Treatment	Dose	Volume of mercury displaced in ml at various time intervals in hours.					
			0	1	2	3	4	5
1	Control	0.1 ml 1% W/V	0.39 ± 0.025	0.49± 0.057	0.62± 0.057	0.65± 0.057	0.67± 0.057	0.69± 0.058
2	Ibuprofen	10 mg/ kg	0.26± 0.005	0.29±*** 0.005	0.33±*** 0.002	0.25±*** 0.003	0.26±*** 0.003	0.19±*** 0.04
3	<i>Tamilnadia uliginosa</i> Methanolic extract	100 mg/ kg	0.39± 0.01	0.47± ^{NS} 0.02	0.57±* 0.05	0.60±* 0.03	0.59±** 0.01	0.57±** 0.04
4	<i>Tamilnadia uliginosa</i> Methanolic extract	200 mg/ kg	0.38± 0.01	0.45±* 0.02	0.53±** 0.05	0.54±** 0.03	0.54±*** 0.01	0.52±*** 0.04
5	<i>Tamilnadia uliginosa</i> Methanolic extract	400 mg/kg	0.37± 0.01	0.43±** 0.02	0.43±*** 0.05	0.31±*** 0.03	0.31±*** 0.01	0.29±*** 0.04

Significance of difference in control and extracts treated groups were determined by one-way analysis of variance (ANOVA) *** P<0.001 **P<0.01, * P<0.05 are significant and ^{NS} P> 0.05, are not significant. All values are means of individual data obtained from six rats (n=6)

Table 2. Percentage inhibition of *Tamilnadia uliginosa* extracts on carrageenan induced paw oedema in rats

S. No	Treatment	Dose	% Inhibition of paw volume at various time intervals					
			0	1	2	3	4	5
1	Control	0.1 ml 1% W/V	00	00	00	00	00	00
2	Ibuprofen	10 mg/ kg	00	40	46	61	61	72
3	<i>Tamilnadia uliginosa</i> Methanolic extract	100 mg/ kg	00	4	8	7	11	17
4	<i>Tamilnadia uliginosa</i> Methanolic extract	200 mg/ kg	00	8	14	16	19	24
5	<i>Tamilnadia uliginosa</i> Methanolic extract	400 mg/ kg	00	12	30	52	53	57

Fig 1. Histogram showing the anti-inflammatory activity of various extracts of *Tamilnadia uliginosa*

DISCUSSION AND CONCLUSION

Conventional NSAIDs of modern medicine are confronted with common and serious unacceptable side effects as well as adverse effects notably gastrointestinal bleeding, hepato nephrotoxicity (Waheed *et al.*, 2011). Carrageenan induced edema is commonly used as an experimental animal model for acute inflammation and is believed to be biphasic, of which the first phase is mediated by the release of histamine and 5-HT followed by kinin release and then prostaglandin in the later phase (Amosora *et al.*, 1998; Alcaraz and Jimenez, 1988). Prostaglandin-E₂ (PGE₂), a powerful vasodilator, synergizes with other inflammatory vasodilators such as histamine, bradykinin and serotonin (Mohamed *et al.*, 2011; Georgewill *et al.*, 2010; Foyet *et al.*, 2011) and contributes to redness as well as increased blood flow in areas of acute inflammation.

REFERENCES

- Alcaraz MJ and Jimenez MI. Flavonoids as anti-inflammatory agents. *Fitoterapia*, 59, 1988, 25-38.
- Amosora KM, Iarenko DB, Potapkv OY, Ivashchenko NA. A comparative evaluation of the efficacy of peroral and intravenous pulse therapy with methylprednisolone in rheumatic diseases. *Lik sprava*, 7, 1998, 48.

The results in the study indicate that pretreatment with the selected plant extracts and ibuprofen suppressed the increased paw oedema produced by the phlogistic agent. Phytochemical screening confirmed the presence of flavonoids, tannins and saponins in the methanolic extract of roots of *Tamilnadia uliginosa*. Flavonoids have been reported to play role in analgesic activity by targeting prostaglandins and are known to inhibit prostaglandin synthetase, more specifically the endoperoxides (Rajnarayana *et al.*, 2001; Rao *et al.*, 1998). The results obtained, scientifically support the folklore pertaining to the medicinal properties of *Tamilnadia uliginosa*.

ACKNOWLEDGEMENTS

The authors were thankful to College of Pharmaceutical Sciences, Andhra University for providing the facilities to carry out the present research work.

- Conforti F, Sosa S, Marrelli M, Menichini F, Statti GA, Uzunov D, et al. The protective ability of Mediterranean dietary plants against the oxidative damage: the role of radical oxygen species in inflammation and the polyphenol, flavonoid and sterol contents. *Food Chemistry*, 112(3), 2009, 587–594.
- Coussens LM, Werb Z. Inflammation and cancer. *Nature*, 420(6917), 2002, 860–67.
- Dharmasiri MG, Jayakody JRAC, Galhena G, Liyanage SSP, Ratnasooriya WD. Anti-inflammatory and analgesic activities of mature fresh leaves of *Vitex negundo*. *Journal of Ethnopharmacology*, 2(3), 2003, 199–206.
- Duffy JC, Dearden JC, Rostron C. Design, Synthesis and biological testing of a novel series of anti-inflammatory drugs. *J. Pharm. Pharmacol*, 53, 2001, 1505-1514.
- Foyet HS, Abdou BA, Ponka R, Asongalem AE, Kamtchouing P, Nastasa V. Effects of *Hibiscus asper* leaves extracts on carrageenan induced oedema and complete Freund's adjuvant-induced arthritis in rats. *Journal of Cell and Animal Biology*, 5, 2011, 69-75.
- Georgewill OA, Georgewill UO, Nwankwoala RNP. Anti-inflammatory effects of *Moringa oleifera* lam extract in rats. *Asian Pac J Trop Med*, 3, 2010, 133-135.
- Georgewill OA, Georgewill UO. Evaluation of the anti-inflammatory activity of extract of *Vernonia amygdalina*. *Asian Pac J Trop Med*, 3, 2010, 150-151.
- Gupta M, Mazumder UK, Gomathi P, Selvan VT. Anti-inflammatory evaluation of leaves of *Plumeria acuminata*. *BMC Complementary and Alternative Medicine*, 6, 2006, 36.
- Jacob Jesurun RS, Senthilkumaran Jagadeesh, Somasundaram Ganesan, Venugopala Rao K, Madhavi Eerike. Anti inflammatory activity of ethanolic extract of *Nymphaea alba* flower in swiss albino mice. *Int J Med Res Health Sci*, 2(3), 2013, 474-478.
- Kumara N. Identification of strategies to improve research on medicinal plants used in Sri Lanka, in *Proceedings of the WHO Symposium*, 12–14, University of Ruhuna, Galle, Sri Lanka, 2001.
- Mohamed STK, Azeem AK, Dilip C, Sankar C, Prasanth NV, Duraisami R. Anti-inflammatory activity of the leaf extracts of *Gendarussa vulgaris* Nees. *Asian Pac J Trop Biomed*, 1, 2011, 147-149.
- Nathan C. Points of control in inflammation. *Nature*, 420(6917), 2002, 846–852.
- Oyedapo OA, Adewunmi CO, Iwalewa EO, Makanju VO. Analgesic, antioxidant and anti-inflammatory related activities of 21-hydroxy-2,41-dimethoxychalcone and 4-hydroxychalcone in mice. *Journal of Biological Sciences*, 8(1), 2008, 131–136.
- Peskar BM. On the synthesis of prostaglandins by human gastric mucosa and its modification by drugs. *Biochimica et Biophysica Acta*, 487(2), 1977, 307–314.
- Rajnarayana K, Reddy MS, Chaluvadi MR, Krishna DR. Biflavonoids classification, pharmacological, biochemical effects and therapeutic potential. *Indian Journal of Pharmacology*, 33, 2001, 2-16.
- Rao MR, Rao YM, Rao AV, Prabhkar MC, Rao CS Muralidhar N. Antinociceptive and anti-inflammatory activity of a flavonoids isolated from *Caralluma attenuate*. *Journal of Ethnopharmacology*, 62, 1998, 63-66.
- Robert A. Antisecretory, antiulcer, cytoprotective and diarrheogenic properties of prostaglandins. *Advances in Prostaglandin and Thromboxane Research*, 2, 1976, 507–520.
- Tapiero H, Nguyen Ba G, Couvreur P, Tew KD. Polyunsaturated fatty acids (PUFA) and eicosanoids in human health and pathologies. *Biomedicine and Pharmacotherapy*, 56(5), 2002, 215–222.
- Waheed MA, Farha D, Rabbani S, Yousuf Ali M, Zeeyaaddin K, Ali AA. Animal studies on unani drug formulation for its Analgesic, Antipyretic, Anti-inflammatory and Non-Toxic effect. *Int J Pharm Pharm Sci*, 3(4), 2011, 66-72.