



CLINICAL UTILITY OF LIPID PROFILE, HOMOCYSTEINE IN EARLY STAGES OF ACUTE MYOCARDIAL INFARCTION

Subramanyam Naidu P¹, Deepthi Vodnala^{2*}, E.Prabhakar Reddy³

¹Assistant Professor of General Medicine, Sri Lakshmi Narayana Institute of Medical Sciences, (Affiliated to Bharath University, Chennai), Pondicherry, India.

²Assistant Professor of General Medicine, Sri Lakshmi Narayana Institute of Medical Sciences, (Affiliated to Bharath University, Chennai), Pondicherry, India.

³Associate Professor of Biochemistry Department & Central lab HOD, Sri Lakshmi Narayana Institute of Medical Sciences, (Affiliated to Bharath University, Chennai), Pondicherry, India

ABSTRACT

Coronary artery disease is now a major health problem in India. In past few decade, the battle to reduce the incidence of CAD has led the researchers to look for various clinical markers, which would help early diagnosis of the diseases. The present study was undertaken to assess the levels of lipid profile like Triacylglycerol level, Cholesterol, LDL and Homocysteine in selected Myocardial infarction (MI) patients. Fasting blood samples were taken from 50 MI patients (30 males and 20 females) and compared with age matched controls. Blood samples from MI patients were taken at Zero hours (i.e; at the time of admission), 1 hour and after 6 hours. By the end of 6 hrs, TGL, LDL and Homocysteine increases whereas cholesterol does not showed significant difference when compared with controls. Thus the present study indicates a strong association between the lipid profile and homocysteine level with the AMI and thus implying possible risk factors for MI.

Key words: AMI, CAD, Cardiac enzymes, LDL, Homocysteine.


Corresponding Author: **Dr. Deepthi Vodnala** Email: drpebyreddy@gmail.com

INTRODUCTION

Acute myocardial infarction (AMI) is the most important consequences of coronary artery disease. CAD is the most common cause of death. It is responsible for the deaths of approximately one in five and one in six women. MI is now considered as a part of spectrum referred to as Acute Coronary Syndrome. Although traditional risk factors of MI are helpful in diagnosis, specific clinical markers would be valuable in identifying the persons who are at risk (Angeliene T, *et al.* 2003). In the UK, the highest recorded rates of CAD mortality are people

born in India, Pakistan, and Bangladesh. South Asians are thought to have a 40-60% higher risk of CHD related mortality compared to other population and this is due to use of ghee in cooking, raised blood glucose, cholesterol, parental consanguinity, paternal history of CVD, low level of income and lack of education (Ismail J, *et al.*). The average incidence of myocardial infarction for those aged between 30 and 69 years is about 600 per 1,00,000 for men and 200 per 1,00,000 for women. Incidence increases with age and elderly people also tend to have higher of morbidity and mortality from their infarcts.

Risk factors may be increasing age, male, family history of premature history of premature CHD, premature menopause. Modifiable risk factors for atherosclerosis include smoking, Diabetes mellitus, metabolic syndrome, hypertension, hyperlipidaemia, obesity and physical inactivity. Diagnosis of infarction can be achieved by ECG changes and elevation of cardiac

Access this article online		
Home page: http://onlineijp.com/	Quick Response code	
		
Received:25.04.2015	Revised:02.06.2015	Accepted:15.07.2015

markers .in addition to help diagnosis of AMI ,the laboratory parameters are of major importance in monitoring cause of infarction and in estimating its size now a days highly playing important role in diagnosis of AMI. (Hitesh shah, *et al.* 2003). The pathogenesis of AMI is multi factorial however ,several studies have implicated impaired lipid metabolism as one of the crucial factors in the development of the disease. (Haseeb A. 2013). So much attention has been focused on serum lipid profile because of their strong association of CAD.

The above literature clearly indicates important role of lipid metabolism in AMI. However, biomarker value of various components of lipid profiles is not clear due to conflicting findings in various studies. We therefore compared the lipid profile of AMI with respect to normal subjects (Haseeb A. 2013). There are some cases with MI those who don't have any of the traditional risk factors .So, attention has been focused on other predisposing factors which may contribute to MI. Researchers have taken effort to find the possible association between Plasma Homocysteine levels in AMI. (Angeliene T. 2005) Several studies conducted in different parts of the world was reported the elevated levels of Plasma Homocysteine are associated with CAD, independent of other risk factors .

We therefore compared the cardiac enzymes, lipid profiles and homocysteine levels in AMI with respect to normal subjects. Therefore this has been taken due to the incidence of mortality in MI is high.

Materials and methods

50 patients out of which 30 males and 20 females admitted to the intensive care unit of Sri Lakshmi Narayana Medical college, Pondicherry with documented AMI showing characteristic ECG signs and rise in troponin I concentrations were selected and compared with 50 age matched controls who were volunteers in our hospital. The patients who were under recent surgery, infections, chronic inflammatory diseases, significant hepatic or renal dysfunction and malignancy were excluded from the study.

Sample collection:

Fasting blood was collected from the patients within 6 hrs from the onset of the infarction..Plasma was separated from collected blood sample using EDTA coated tubes. Serum was separated by centrifugation at 2000 rpm and is used for the estimation of lipid profile such as total cholesterol, TG's, and LDL. The separated plasma was used for analyzing the plasma homocysteine using ELISA.

The values of lipid profile and homocysteine levels in patients of AMI were compared with the normal subjects.

Result

Myocardial infarction can occur at any age irrespective of sex, but generally it is a disease of middle and old age. The result of the present study is given as below.

Table 1: Lipid profile and Homocysteine

Parameters	Males (30)	Female (20)	First hour	After 6 Hrs	Controls	P value
Cholesterol	177.07±8.49	183±8.75	-	-	188.5±8.38	<0.001
TG	131.23±12.54	139.26±12.81	-	-	117.7±12.34	<0.001
LDL	143.4±6.84	153.1±6.83	-	-	115.03±2.38	<0.001
Homocysteine	16.56±1.12	18.91±1.83	-	-	12.40±0.99	<0.001

DISCUSSION

The diagnosis of acute myocardial infarction (AMI) as traditionally been based on the characteristic clinical history, ECG abnormalities and increased serum concentrations of cardiac markers. In case of lipid profile, our results showed significant increase in TG and LDL levels in AMI patients but the cholesterol level shows no significant raise as compared with the controls. The mechanism of increase in TG's after acute MI may be due to elevated flux of fattyacids and impaired removal of VLDL from the plasma. (Nigam P.K, *et al.* 2004) Another possible mechanism for elevated TG levels may be the effect of β blockers but this seems to be invalid for increased TG levels on day 3 and pre-discharge (day 7) as β blockers take about two weeks to show their effect on serum lipids. (Nigam P.K, *et al.* 2004). LDL cholesterol recorded high significant changes throughout the study period in our subjects. However a significant decrease or

no significant change in LDL cholesterol following MI has been reported by others. (Prasad K. 1999). We found no significant changes in serum total cholesterol levels either during the AMI or after 3 months. Our study provides information which is direct contrast to that by others who found either a decrease or increase during the acute phase of MI. (Nigam P.K, *et al.* 2004, Ueland, P.M, *et al.* 1993)

There is an elevation of Plasma Homocysteine in AMI patients when compared with control group (table 2). Statistical significant difference were observed in the mean of Plasma Homocysteine concentrations between the AMI patients (male 16.56±1.12 mM/L, females 18.91±1.83) and in controls (12.40± 0.99mM/L).[6] .Recent report on Homocysteine suggests that it is a independent predictor of vascular disease including stroke and coronary artery disease. Mutations in the methyl tetrahydrofolate reductase gene are one of the most

frequent causes of moderately elevated plasma Homocysteine. (Haseeb A. 2013)

Hyperhomocysteinemia may also occur due to nutrition deficiency that leads to low blood concentration of folate, Vit B₁₂ or vitamin B₆.

Homocysteine may also induce atherosclerosis by effecting endothelial derived relaxin factor Nitric oxide(NO). It also enhances the lipid peroxidation which may decrease the expression of endothelial NO synthase and directly degrade NO. The finding of this study

underscores the importance of determining the levels of Plasma Homocysteine in individuals who are at risk of developing AMI (Ismail J, et al)

CONCLUSION

In Conclusion , our data provides a clear evidence that serum enzymes, lipid profiles and Homocysteine levels are markedly useful in determining the patients of Acute Myocardial Infarction.

REFERENCES

- Acute myocardial infarction. www.doctor AMI/Doctor/Patient.co.uk
- Angeliene T, Rita Mary Aruna.K.Ramadevi,G.Mohan,Nirmala Jeyaraj .Serum lipoprotein (a) and lipid profile in young south indian patients with myocardial infarction. *Indian Journal of Clinical Biochemistry*, 2003,18 (1) 103-106.
- Ismail J, TH.Jafar and N Chaturvedi.Risk factors for non fatal myocardial infarction in young south asian ADULTS. www.ncbi.nlm.nih.gov/PMC1768096
- Hitesh shah and N.Haridas. evaluation of clinical utility of serum enzymes and troponin-t in the arly stages of acute myocardial infarction. *Indian Journal of Clinical Biochemistry*, 2003 18(2)93-101.
- Hitesh Shah and N. Haridas. a serial follow up study of cardiac marker enzymes during the week after acute myocardial infarction. *Indian Journal of Clinical Biochemistry*, 2007 / 22 (1) 33-36
- Haseeb A. Khan1, Abdullah S. Alhomida1 and Samia H. Sobki. Lipid profile of patients with acute myocardial infarction and its correlation with systemic inflammation Biomarker Insights 2013:8 1-7
- Angeliene T, Rita Mary Aruna.K.Ramadevi,G.Mohan,Nirmala Jeyaraj. Homocysteine status and acute myocardial infarction among tamilians. *INDIAN JOURNAL OF CLINICAL BIOCHEMISTRY*, 2005,20(1) 18-20
- Nigam P.K, V.S. Narain and M.Hasan. Serum lipid profile in patients with acute myocardial infarction. *Indian Journal of Clinical Biochemistry*,2004,19 (1) 67-70
- Myocardial inarction-wikipwedia
- Prasad ,K. (1999) Homocysteine, a risk factor for cardiovascular disease. *Int.J of Angiol*.8,76-86.
- Ueland, P.M.,Refsum,H., Stabler,S.P.,Malinow,M.R., Andersson,A., Allen,R.H.(1993).Total homocysteine in plasma or serum: methods and clinical applications. *Clin Chem*.39:1764-1779
- A.M Huggon,J Chambers, N Nayeem,P tutt,M C rook,S Swaminathan.Biochemical markers in the management of suspected acute myocardial infarction in the emergency department.2014,15-19..